

(19)

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 769 064 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
28.08.2002 Bulletin 2002/35

(21) Application number: 96912316.5

(22) Date of filing: 27.04.1996

(51) Int Cl.7: **C12P 17/02**

(86) International application number:
PCT/KR96/00060

(87) International publication number:
WO 96/034110 (31.10.1996 Gazette 1996/48)

(54) A METHOD FOR MASS PRODUCTION OF TAXOL BY SEMI-CONTINUOUS CULTURE

METHODE ZUR MASSENPRODUKTION VON TAXOL MITTELS SEMI-KONTINUIERLICHER
KULTUR

PROCEDE DE PRODUCTION DE MASSE DE TAXOL PAR CULTURE SEMI-CONTINUE

(84) Designated Contracting States:
DE FR IT

(30) Priority: 27.04.1995 KR 9510204

(43) Date of publication of application:
23.04.1997 Bulletin 1997/17

(60) Divisional application:
01123786.4 / 1 164 197

(73) Proprietor: **Samyang Genex Co., Ltd.**
Chongno-Ku, Seoul 110-470 (KR)

(72) Inventors:
• **CHOI, Hyung-Kyoon**
Seo-ku, Taejon 302-150 (KR)
• **ADAMS, Tom Lee**
Millbrae, CA 94030 (US)
• **STAHLHUT, Roy William**
Belmont, CA 94002 (US)
• **KIM, Sang-Ic**
Nowon-ku, Seoul 139-220 (KR)
• **YUN, Jeong-Hwan**
Taejon 305-348 (KR)

- **SONG, Bong-Kyu**
Yusong-ku, Taejon 305-390 (KR)
- **KIM, Jin-Hyun**
Taejon 302-222 (KR)
- **SONG, Jun-Seog**
Yusong-ku, Taejon 305-390 (KR)
- **HONG, Seung-Suh**
Yusong-ku, Taejon 305-390 (KR)
- **LEE, Hyun-Soo**
Yangchun-ku, Seoul 158-070 (KR)

(74) Representative:
TER MEER STEINMEISTER & PARTNER GbR
Patentanwälte,
Mauerkircherstrasse 45
81679 München (DE)

(56) References cited:
WO-A-93/17121 **WO-A-93/19585**
US-A- 5 019 504

Remarks:

The file contains technical information submitted
after the application was filed and not included in this
specification

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 769 064 B1

Description

BACKGROUND OF THE INVENTION5 Field of the Invention

[0001] The present invention relates to a method for mass production of taxol from Taxus genus plant cell culture, more specifically, to a method for mass production of taxol by semi-continuous culture of Taxus genus plant cell with a high yield.

10

Description of the Prior Art

[0002] Taxanes are diterpene compounds containing a taxane skeleton. For example, taxol is famous as the first identified compound with a taxane ring which was isolated from the bark of the pacific yew, Taxus brevifolia, which is effective for the treatment of leukemia and cancer. Recently, it has been reported that: taxol is capable of curing approximately 30%, 50% and 20% of ovarian, breast and lung cancer patients, respectively, by inhibiting depolymerization of microtubules(see: E.K. Rowinsky et al., J. Natl. Cancer. Inst., 82:1247-1259(1990)).

15

[0003] On the other hand, total chemical synthesis, semi-synthesis and extraction methods have been employed to prepare taxol.

20

[0004] The total chemical synthesis method, however, has not been practically applied in the art, since it requires very expensive chemical reagents and the yield is not so high, which can be expected from the complicated chemical structure of taxol.

25

[0005] The semi-synthesis method employing precursors such as 10-deacetylbaccatin III, has revealed some drawbacks since it essentially entails complicated and multiple steps of isolating and purifying the taxol precursors from Taxus genus plant and transforming the precursors to taxol.

30

[0006] In this regard, the extraction method by which taxol can be isolated from Taxus genus plants in a direct manner, has prevailed in the art, since it has the advantage of economy. However, the said method has proven to be less satisfactory in the sense that it essentially requires a large amount of yew trees to purify taxol, which finally gives rise to the serious environmental disruption.

35

[0007] Accordingly, the ability of total chemical synthesis, semi-synthesis and extraction method to supply taxol for world-wide chemotherapeutic use is not assured; and, there are strong reasons for exploring and developing alternative means of taxol production.

40

[0008] As a promising alternative to solve said problems, the cell culture method for taxol production has been proposed in the art.

45

[0009] The cell culture-based process for taxol production, unlike the prior art, has the following advantages as followings: first, taxol can be produced in a steady manner, regardless of fluctuation of supply of yew plants due to the damage by blight and harmful insects, etc.; secondly, cell cultures can be propagated in large bioreactors, from which taxol can be massively produced by manipulating culture conditions; thirdly, cell cultures produce a simpler spectrum of compounds compared to other prior art methods, considerably simplifying separation and purification; fourthly, a cell culture process can adapt quickly to rapid changes in demand better than the other methods; fifthly, a cell culture process can produce taxol as well as taxane precursors such as baccatin that can be converted to taxol.

50

[0010] Methods for producing taxol by utilizing cultured plant cells have been described in the art:

55

[0011] USP 5,019,504 discloses a method for producing taxol and its derivative utilizing cultured cells of Taxus brevifolia. However, the yield of taxol described therein is 1-3 mg/L, which is insufficient for industrial application. Besides, the production of taxol by the cell culture is unstable and even when a primary cell of high productivity can be obtained by selection, it is difficult to keep its content by subculturing(see: E. R. M. Wickremesine et al., World Congress on Cell and Tissue Culture(1992)).

60

[0012] USP 5,015,744 teaches a semi-synthetic method from baccatin III, which is a precursor in biosynthesis of taxol. By the use of the plant tissue culture, a raw material for the semi-synthetic process such as baccatin III can be produced, thus the plant tissue culture can also be utilized for taxol production by the above-mentioned semi-synthetic process.

65

[0013] WO 93/17121 offers a method for taxol production by cell culture of Taxus genus plant while changing composition of medium, growth rate, and production rate, etc. In case of Taxus chinensis, 24.1mg/L of taxol can be obtained in 18 days of culture and biomass doubles every 2.5 days.

70

[0014] All of these patents describes methods for mass production of taxol by employing batch culture; there is no teaching in said patents on, nor is there anticipated, semi-continuous culture of a taxol-producing cell line.

[0015] Under the circumstances, USP 5,407,816 describes that Taxus chinensis cells are inoculated to a nutrient medium to form a suspension which is, in turn, cultivated to form a suspension culture which is, in turn, subcultured in

the other nutrient medium to form a producing culture, and which finally gives taxol and taxanes in a yield of 153 mg/L. The said method has considerably improved the productivity of taxol, however, it has been proven less satisfactory in the sense that it essentially requires so many nutrient media whose compositions are so complicated, and high productivity can be realized under rather a limited growth condition.

5 [0016] Therefore, there is a continued need to develop a practical and simple method for taxol production, which is able to meet the requirement of high productivity which is a critical factor to determine whether it can be utilized in industrial applications or not.

10 SUMMARY OF THE INVENTION

[0017] In accordance with the present invention, it has been discovered that taxol can be produced with a high degree of efficiency, by the semi-continuous culture of a novel taxol-producing cell line, Taxus chinensis SYG-1.

[0018] A primary object of the present invention is, therefore, to provide a method for mass production of taxol by semi-continuous culture.

15 [0019] The other object of the invention is to provide a novel taxol-producing cell line isolated from Taxus chinensis.

[0020] The above primary object is achieved by a method according to claim 1.

BRIEF DESCRIPTION OF THE DRAWINGS

20 [0021] The above and the other objects and features of the present invention will become apparent from the following descriptions given in conjunction with the accompanying drawings, in which:

Figure 1 is a graph showing the effect of AgNO_3 on the taxol productivity.

25 Figure 2 is a graph showing the effect of NH_4 -citrate on the taxol productivity.

Figure 3 is a graph showing the effect of maltose on the taxol productivity.

30 Figure 4 is a graph showing the productivity of taxol in semi-continuous culture of SYG-1 in accordance with the culture cycle.

DETAILED DESCRIPTION OF THE INVENTION

35 [0022] The present inventors first developed a taxol-producing cell line, based on the callus induced from Taxus chinensis, and compared the cell line with those known in the art. From the comparative studies on the morphological, physiological and growth condition of the taxol-producing cell lines, it has been determined that the newly developed cell line is a novel one which is somewhat different from those of prior art, in light of taxol productivity, mode of taxol secretion, etc. Accordingly, the cell line was named Taxus chinensis SYG-1 (hereinafter, referred to as "SYG-1" for convenience) and it was deposited with an International Depositary Authority (IDA), the Korean collection for Type Cultures (KCTC), Korea Research Institute of Bioscience and Biotechnology, on March 14, 1996, under an accession number of KCTC 0232BP.

40 [0023] Then, the inventors optimized the growth condition of the cell line of the invention, and determined that: SYG-1 cells grown well on B5 medium, most preferably B5 medium supplemented with 20 μM NAA (naphthoxyacetic acid), 0.4 μM BAP (6-benzylaminopurine), 1g/L casein hydrolysate and 30g/L sucrose, under a condition of a temperature of 24°C, at an agitation speed of 150rpm. On the other hand, effects of AgNO_3 , NH_4 -citrate and maltose on the productivity of taxol were also examined, and the inventors concluded that their addition considerably improved the productivity of taxol.

45 [0024] According to the present invention, taxol can be prepared with a high yield, by employing semi-continuous culture of SYG-1, which comprises:

- 50
- (i) inoculating Taxus genus plant cell on a medium containing 1 to 10% (w/v) sugar, and incubating it; and,
 - (ii) transferring 1/10 to 1/2 volume of the culture obtained in the step (i) to a fresh medium and repeating the step (i), adding 1 to 10% (w/v) sugar to the remnant culture and incubating to the time of maximum production of taxol.

55 [0025] At the beginning of culture, AgNO_3 may be added to Taxus genus plant cell culture which is incubated for 10 to 20 days, more preferably 10 to 15 days, at a concentration of 1 to 15 μM , more preferably 5 to 10 μM . Furthermore, NH_4 -citrate and maltose may be added to the culture after 5 to 30 days, more preferably 5 to 10 days of incubation, at a concentration of 1 to 15mM, more preferably 1 to 10mM and 1 to 10% (w/v), more preferably 1 to 5% (w/v), respec-

tively.

[0026] After 5 to 30 days of incubation, 1/10 to 1/2 volume of total culture is transferred to other flask containing fresh medium whose contents are the same as the one employed at the beginning of the culture, and started a cycle of culture. Then, remnant culture equivalent to 9/10 to 1/2 volume of total culture, was added 1 to 10%(w/v), more preferably 1 to 5%(w/v) of maltose, and incubated for 30 to 60 days in which taxol production is maximized. At this time, incubation was made at 24°C, agitation speed of 150rpm under a dark condition.

[0027] Taxus genus plant employed in the method of the present invention includes Taxus brevifolia, Taxus canadensis, Taxus cuspidata, Taxus baccata, Taxus globosa, Taxus floridana, Taxus wallichiana, Taxus media and Taxus chinensis, however, Taxus chinensis SYG-1 which is newly developed by the inventors, is most preferably employed therein.

[0028] According to the method of present invention, the productivity of taxol to the level of 300mg/L, after 42 to 49 days incubation, is observed, when Taxus chinensis SYG-1 is employed in the semi-continuous culture with the addition of AgNO₃, NH₄-citrate and maltose to improve the productivity of taxol.

Quantitative Analysis of Taxol

[0029] Taxol which is produced from the culture of Taxus chinensis SYG-1 according to the method of the present invention, is quantitatively assayed by employing high performance liquid chromatography under a specific condition described in Table 1 below.

Table 1:

Condition for quantitative assay of taxol	
Instrument	HPLC(Waters, U.S.A.)
Column	Capcell Pack C ₁₈ UG 120 (length: 250mm, inner diameter: 4.6mm).
Column temp.	40°C
Mobile phase	CH ₃ CN : water(20 ~ 100% gradient)
Fluid speed	1.0 ml/min
Injection volume	10 µl
Detector	UV(227nm), ATTE = 3

[0030] The present invention is further illustrated in the following examples, which should not be taken to limit the scope of the invention.

Example 1: Induction and characterization of callus from Taxus chinensis

[0031] The cell line of the present invention was isolated from Taxus chinensis by way of callus induction. When plant tissue was incubated on a proper medium containing a hormone, undifferentiated cell of callus was induced: Tissues from bark, needles, stems and roots of Taxus chinensis were washed with tap water and sterilized with Ca-hypochloride solution for 20 to 30min. Then, the sterilized tissue was washed with distilled water 2 to 3 times, chopped to have a length of 1cm, and transferred to B5 medium(see: Gamborg et al., Can. J. Biochem., 46:417-421(1968)) supplemented with 20µM NAA(naphthoxyacetic acid), 0.4µM BAP(6-benzylaminopurine), 1g/L casein hydrolysate and 30g/L sucrose after solidifying with 0.2%(w/w) gelite, and incubated at a temperature of 24 to 26°C for 2 to 6 weeks under a dark condition to induce the callus of interest.

[0032] The growth medium for callus was selected by transferring the callus to several media well known in the art, e.g., MS(see: Murashige, T. and F. Skoog, F., Physiol. Plant, 5:473(1962)), SH(see: Schenk and Hilderbrandt, Can. J. Bot., 50:199-204(1972)), WPM(see: Lloyd and Mccown, Int. Plant Prop. Soc. Proc., 30:421-427(1981)) and B5 media (see: supra), respectively, and observing the growth pattern of the callus by naked eye. From the results above, it was determined that the callus showed good growth on B5 medium containing 20µM NAA, 0.4µM BAP and 2g/L casein hydrolysate.

[0033] Then, studies on the morphology, physiology and growth condition were made for the cell line thus induced, and compared with those of known in the art; and, the results were summarized in Tables 2 and 3 below. As can be seen in Tables 2 and 3, it was determined that the cell line which has been developed in the invention has a distinction over those of prior art, in light of taxol productivity and mode of taxol secretion, etc. Accordingly, the cell line was named Taxus chinensis SYG-1 and was deposited with an International Depositary Authority(IDA), the Korean Collection for

Type Cultures(KCTC) on March 14, 1996, under an accession number of KCTC 0232BP.

Table 2:

Morphology and growth condition of <u>Taxus chinensis</u> SYG-1	
<u>Taxus chinensis</u> SYG-1	Characteristics
Size	50-150μm
Mobility	(-)
Agglutination	Weak
Aggregation	(+)
Adaptation to shear stress	Strong
Color of culture	Weak brown
Culture temperature	24°C.

Table 3: Comparison of characteristics of cell lines of the invention and prior art

Characteristics	SYG-1	USP 5407816	USP 5019504	WO 92/13961
Secretion mode of taxol	intracellular	extracellular	not clear	not clear
Light-requirement	dark	light	dark	dark
Taxol productivity	~300mg/L	153mg/L	1-3mg/L	0.04% DCW

Example 2: Suspension culture of SYG-1

[0034] To select the most preferred medium for the suspension culture of SYG-1, the cell line was transferred to MS (Murashige and Skoog medium), SH (Schenk and Hildebrandt medium), WPM (Woody Plant Medium) and B5 (Gamborg's B5 medium) media, respectively, and the growth pattern was observed analogously in Example 1, which resulted that SYG-1 cell was well grown on B5 medium containing 20 μ M NAA, 0.4 μ M BAP and 2g/L casein hydrolysate, like the callus. In addition, it was also determined that SYG-1 cell was well grown at a temperature of 24°C, at an agitation speed of 150rpm.

[0035] A serial cultivation of the callus was made every 4 weeks in a case of solidified medium, while transferring a part of tissue by the aid of forceps: A piece of the callus was maintained on the plate of solid medium. Then, the callus maintained on the solid medium was inoculated into a small volume of B5 medium, and small amounts of medium were supplemented to the culture, as the cells grew to increase total volume of culture. Then, well grown cell lines were diluted in the modified B5 medium contained in 500ml Erlenmeyer flask at a ratio of 1/5(v/v), and inoculated into suspension culture medium every two weeks.

Example 3: Effect of AgNO₃ on the taxol productivity

[0036] It has been well known that AgNO₃ is an antagonist of the plant hormone, ethylene, which effects on the plant cell growth and secondary metabolite production. Accordingly, the present inventors tested the effect of AgNO₃ on the taxol production in a suspension culture of SYG-1.

[0037] To a 250ml of Erlenmeyer flask, was poured 75ml of B5 medium containing 10 μ M AgNO₃ and 25ml of 14 day old cell culture was inoculated into this medium, and incubated in an analogous manner described in Example 2. Then, the productivity of taxol was compared with a control which does not contain AgNO₃(see: Fig. 1). As can be seen in Fig. 1, it was clearly determined that the amount of taxol produced, 98.95mg/L when AgNO₃ was added to a medium (-●-), was 4.7 times the control (-○-).

Example 4: Effect of NH₄-citrate on the taxol productivity

[0038] The productivity of taxol in a suspension culture of SYG-1 was monitored, after the addition of NH₄-citrate. To a 250ml of Erlenmeyer flask was poured 75ml of B5 medium and 25ml of 14 day old cell culture was inoculated into this medium, and incubated at the same growth condition described in Example 2. Then, 5mM of NH₄-citrate was added to the culture after 9 days of incubation, and the taxol productivity was compared with a control in which NH₄-citrate was not added(see: Fig. 2). As can be seen in Fig 2, it was determined that the amount of taxol produced was about 103.6mg/L when NH₄-citrate was added to a medium (-●-), which was 4.9 times the control(-○-).

Example 5: Effect of maltose on the taxol productivity

[0039] It has been well known that the increase of sugar concentration in a plant cell culture leads to an increase in secondary metabolite production. For example, it has been reported that: addition of 3%(w/v) sucrose and 5%(w/v) mannitol has improved the productivity of antocyanin in callus culture of *Daucus carota* (see: Knobloch, K. -H., et al., Zeitschrift für Naturforschung, 35c:55-556(1981)); and, 88mM sucrose and 165mM mannitol heightened the antocyanin productivity in a suspension culture of *Vitis vinifera* (see: Rajendran, L., et al., Biotechnology Letters, 14(8):707-712 (1992)). On the other hand, the decrease in the growth and the secondary metabolite production were also reported when the amount of sugar over a certain concentration was added, possibly due to osmotic stress(see: Do, C. B., and Cormier, F., Plant Cell Reports, 9:500-504(1990)).

[0040] The effect of maltose addition to SYG-1 was evaluated.

[0041] To a 250ml of Erlenmeyer flask was poured 75ml of B5 medium and 25ml of 14 day old cell culture was inoculated into this medium, and incubated under the same growth condition described in Example 2. Then, 2%(w/v) and 3%(w/v) of maltose were added to the culture after 9 days and 21 days of incubation, respectively and the productivity of taxol was compared with a control which does not contain maltose(see: Fig. 3). As can be seen in Fig. 3, it was determined that the amount of taxol was 112.75mg/L when maltose was added to a medium(-●-), which was 5.3 times the control (-○-).

Example 6: Semi-continuous culture of SYG-1

[0042] Based on the results obtained in Examples 2 to 5, taxol was prepared by employing semi-continuous culture of SYG-1.

[0043] To a 250ml of Erlenmeyer flask was added 80ml of growth medium and 20ml of 14 day old SYG-1 culture,

10µM AgNO₃ was added at the beginning of culture, followed by the addition of 5mM NH₄-citrate and 2%(w/v) maltose after 9 days of incubation. After 21 days of incubation, 20ml of culture equivalent to 1/5 volume of total culture was transferred to other flask containing 80ml fresh medium to start another cycle of culture. Then, to 80ml of remnant culture equivalent to 4/5 volume of total culture, was added 3%(w/v) maltose, and incubated for additional 21 days in which taxol production is maximized. At this time, incubation was made at 24°C, and an agitation speed of 150rpm. Culture samples were taken, periodically to check for microbial contamination. The amount of taxol produced in the culture was also determined by employing high performance liquid chromatography(HPLC, Waters, U.S.A.) described as above. As a result, microbial contamination was not observed and the productivity of taxol after 42 days incubation was determined to be 284mg/L(see: Fig. 4). As can be seen in Fig. 4, taxol concentration was increased as the cycles of culture were repeated.

[0044] As clearly illustrated and demonstrated as above, the present invention provides a method for mass production of taxol by semi-continuous culture of Taxus genus plant with a high yield. According to the present invention, the productivity of taxol to the level of 300mg/L, after 40 to 50 days incubation, was observed, when Taxus chinensis SYG-1 was employed in the semi-continuous culture with the addition of AgNO₃, NH₄-citrate and maltose to improve the productivity of taxol.

Claims

1. A method for mass production of taxol by semi-continuous culture, said method comprising the steps of:
 - (i) inoculating Taxus genus plant cells in a medium containing 1 to 10% (w/v) sugar and incubating it,
 - (ii) transferring a part of the culture obtained in step (i) to a fresh medium, such that the volume ratio of said culture to said fresh medium is in the range of 1/10 to 1/2, and repeating step (i), and
 - (iii) adding sugar to the remnant culture at a concentration of 1 to 10% (w/v) and incubating it to the time of maximum production of taxol.
2. The method of claim 1, wherein the Taxus genus plant is selected from the group consisting of Taxus brevifolia, Taxus canadensis, Taxus cuspidata, Taxus baccata, Taxus globosa, Taxus floridana, Taxus wallichiana, Taxus media and Taxus chinensis.
3. The method of claim 1, wherein the Taxus genus plant cell is Taxus chinensis SYG-1 (KCTC 0232BP).
4. The method of claim 1, which further comprises a step of addition of AgNO₃ to the medium at a concentration of 1 to 15 µM at the beginning of cell culture.
5. The method of claim 4, which further comprises a step of addition of NH₄-citrate or maltose, or both, to the medium at a concentration of 1 to 15 mM and 1 to 10% (w/v), after 5 to 30 days of incubation, respectively.
6. The method of claim 1, wherein the time of maximum production of taxol is 40 to 50 days from the beginning of cell culture.

Patentansprüche

1. Ein Verfahren zur Massenherstellung von Taxol mittels semi-kontinuierlicher Kulturen, wobei genanntes Verfahren die Schritte umfasst:
 - (i) Animpfen von Pflanzenzellen der Gattung Taxus in ein Medium, welches 1 bis 10 Gew.-% Zucker enthält und Inkubation davon, und
 - (ii) Überimpfen eines Teiles der Kultur, welche in Schritt (i) erhalten wurde, in ein frisches Medium, so dass das Volumenverhältnis der genannten Kultur zum genannten frischen Medium in dem Bereich von 1:10 bis 1:2 liegt, und Wiederholung von Schritt (1), und
 - (iii) Zufügen von Zucker zum Rest der Kultur auf eine Konzentration von 1 bis 10 Gew.-% und Inkubation bis

zum Zeitpunkt der maximalen Produktion von Taxol.

2. Das Verfahren nach Anspruch 1, worin die Pflanze der Gattung Taxus ausgewählt ist aus der Gruppe umfassend Taxus brevifolia, Taxus canadensis, Taxus cuspidata, Taxus baccata, Taxus globosa, Taxus floridana, Taxus wallichiana, Taxus media und Taxus chinensis.
3. Das Verfahren nach Anspruch 1, worin die Pflanzenzelle der Gattung Taxus chinensis SYG-1 (KCTC 0232BP) ist.
4. Das Verfahren nach Anspruch 1, welches weiterhin einen Schritt der Zugabe von AgNO_3 zu dem Medium, auf eine Konzentration von 1 bis 15 μM zu Beginn der Zellkultur umfasst.
5. Das Verfahren nach Anspruch 4, welches weiterhin einen Schritt der Zugabe von NH_4 -Citrat oder Maltose oder beidem zu dem Medium, auf eine Konzentration von 1 bis 15 mM und 1 bis 10 Gew.-% respektive nach 5 bis 30 Tagen Inkubation umfasst.
6. Das Verfahren nach Anspruch 1, wobei der Zeitpunkt der maximalen Produktion von Taxol 40 bis 50 Tage nach Beginn der Zellkultur beträgt.

Revendications

1. Procédé de fabrication en masse de taxol par culture semi-continue, ledit procédé comprenant les étapes consistant à :
 - (i) inoculer des cellules de plante du genre Taxus dans un milieu contenant 1 à 10 % (p/v) de sucre et à le mettre en incubation,
 - (ii) transférer une partie de la culture obtenue à l'étape (i) dans un milieu frais, de sorte que le rapport de volumes de ladite culture par rapport au milieu frais soit dans l'intervalle de 1/10 à 1/2 et à répéter l'étape (i), et
 - (iii) ajouter du sucre à la culture restante à une concentration de 1 à 10% (p/v) et à la mettre en incubation jusqu'à la durée de production maximale de taxol.
2. Procédé selon la revendication 1, caractérisé en ce que la plante du genre Taxus est choisie dans le groupe constitué de Taxus brevifolia, Taxus canadensis, Taxus cuspidata, Taxus baccata, Taxus globosa, Taxus floridana, Taxus wallichiana, Taxus media et de Taxus chinensis.
3. Procédé selon la revendication 1, caractérisé en ce que la cellule de plante du genre Taxus est Taxus chinensis SYG-1 (KCTC 0232BP).
4. Procédé selon la revendication 1, caractérisé en outre par une étape consistant à ajouter de l' AgNO_3 au milieu à une concentration de 1 à 15 μM au début de la culture des cellules.
5. Procédé selon la revendication 4, caractérisé en outre par une étape consistant à ajouter du citrate de NH_4 ou du maltose, ou les deux, au milieu à une concentration de 1 à 15 mM et 1 à 10% (p/v), après 5 à 30 jours d'incubation, respectivement.
6. Procédé selon la revendication 1, caractérisé en ce que la durée de production maximale de taxol est de 40 à 50 jours depuis le début de la culture des cellules.




FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4